arXiv:math/0403295v3 [math.GT] 1 May 2008

Geodesic laminations and continued fractions

Igor Nikolaev *

Abstract

A topological invariant of a special class of the geodesic laminations on the modular surfaces is constructed. The invariant has a continuous part (the tail of a continued fraction) and a discrete part (the singularity data). It is shown that the invariant is complete, i.e. the geodesic lamination can be recovered from the invariant. The continuous part of the invariant has the geometric meaning of a slope of the lamination on the modular surface.

Key words and phrases: modular surface, geodesic lamination

AMS (MOS) Subj. Class.: 57M50

1 Introduction

The geodesic lamination is a closed set of the disjoint simple geodesics on a Riemann surface S. It is well known that a simple geodesic is either (i) a closed geodesic, (ii) a non-closed spiral geodesic tending to a closed geodesic or (iii) a non-closed geodesic whose limit set is a perfect (Cantor) subset of S [5]. We shall focus on the geodesic laminations, whose leaves are of type (iii).

The Chabauty topology turns the set of such laminations into an important topological space $\Lambda = \Lambda(S)$. For example, the boundary of the Teichmüller space of a surface S is homeomorphic to Λ, except at the points $\lambda \in \Lambda$ corresponding to the laminations with more than one independent ergodic measure [6].

The Λ is a compact Hausdorff topological space. It has a metric measuring the angles between the asymptotic direction (a slope) of the laminations on the surface S. In the simplest case $S = T^2$, all possible slopes are exhausted by the irrationals $\theta \in [0, 2\pi[$ and therefore the boundary of the Teichmüller space of the torus is a unit circle.

In 1936 A. Weil asked about an extension of the classical Poincaré rotation numbers (the slopes) of the flows on the torus to the case of the higher genus surfaces [17]. It was conjectured that passing to the hyperbolic plane might be critical to a solution of the problem, *ibid.* An excellent survey of Anosov [1] gives an account of the Weil's problem after 1936. There are many interesting developments towards a solution of the problem due to Aranson-Grines-Zhuzhoma

---

*Partially supported by NSERC.

[2], Schwartzman [13], Series [14], Moeckel [10] and others, see [1]. Let us mention a spiritually close and influential work of E. Artin [3].

In this note to every lamination (of a fixed class) a slope – an irrational number, θ, is assigned. Technically, the slope θ is defined via a regular continued fraction and the linear transformations of the hyperbolic plane. The slope measures an asymptotic direction of the lamination on the surface. Recall that the continued fractions can be used to study the action of the ordered sequences of the Möbius transformations on the points of the hyperbolic plane $\mathbb{H}$. Namely, if an infinite sequence $\tau_n \in SL(2,\mathbb{C})$ acts on $\mathbb{H}$, then the image of point $z=\infty$ converges to a complex number:

$$\theta = \prod_{n=0}^{\infty} \tau_n(\infty) = a_0 + \cfrac{b_0}{a_1 + \cfrac{b_1}{a_2 + \dots}}, \tag{1}$$

where $a_n, b_n \in \mathbb{C}$ are uniquely defined upon τ_n, see Wall [16], p. 13.

Let λ^* be a singleton, i.e. a geodesic lamination consisting of one simple closed geodesic. The singletons are dense in the space Λ, see e.g. [4], Lemma 4.2.15. We shall use this fact to approximate the elements $\lambda \in \Lambda$ by the sequences of the singletons $\{\lambda_n^*\}$. Namely, recall that each closed geodesic is the axis of an isometry of the hyperbolic plane, cf. §4.2.3 of the appendix. Therefore, the sequences of the closed geodesics $\{\lambda_n^*\}$ are bijective with the sequences of the transformations $\{\tau_n\}$, such that $|tr\ \tau_n| > 2$. It is easy to see that if $\lambda = \lim \lambda_n^*$ is a lamination defined by the singletons, then the slope $\theta = \theta(\lambda)$ of λ can be introduced via the formula (1) with the τ_n as specified above.

Let $\Gamma(N)$ and $\Gamma_0(N)$ be the principal congruence and the Hecke subgroups of the modular group, respectively. The Riemann surface $X(N) = \Gamma(N)\backslash\mathbb{H}$ ($X_0(N) = \Gamma_0(N)\backslash\mathbb{H}$) is known as a modular surface. Further we deal with the geodesic laminations on the modular surfaces. Since $\tau_n \in SL(2,\mathbb{Z})$, the slope of a $\lambda \in \Lambda(X(N))$ (or $\Lambda(X_0(N))$) can be defined by a regular continued fraction.

To recover the lamination from a slope, one needs to specify the number and type of the boundary components of the lamination, see [5]. Let $g = g(N)$ be the genus of the modular surface and $\Delta = (k_1, \dots, k_m)$ a finite set of the positive integers and half-integers such that $\sum k_i = 2g-2$. The Δ is called a *singularity data* of the lamination. We consider a class of the laminations, which we shall call the *Legendre laminations*, and prove the following theorem.

Theorem 1 *There exists a one-to-one correspondence between the equivalence classes of the Legendre laminations $\lambda \in \Lambda(X(N))$ (or $\Lambda(X_0(N))$) and the pairs (Θ, Δ), where $\Theta = \theta \bmod GL(2,\mathbb{Z})$ is an equivalence class of the positive irrationals modulo the action of the matrix group $GL(2,\mathbb{Z})$ and Δ is the singularity data of the lamination λ.*

The article is organized as follows. In section 2 the geodesic laminations are reviewed. In section 3 the Legendre laminations are introduced and their slopes are defined. In the same section, theorem 1 is proved. The section 4 is reserved for a reference on the continued fractions and the modular surfaces.

Acknowledgements. I wish to thank A. Marden, K. Rafi, M. Sakuma and D. Silver for an interest in the work.

2 Geodesic laminations

This section is a brief review of the geodesic laminations following [5] and contains no original results. We refer the reader to the above cited monograph for a detailed account.

2.1 The Chabauty topology

Let S be a connected complete hyperbolic surface. By a *geodesic* on S we understand the maximal arc consisting of the locally shortest sub-arcs. The geodesic is called *simple* if it has no self-crossing or self-tangent points. We shall leave aside a non-trivial question of the existence of simple geodesics, referring the reader to [11]. A *geodesic lamination* on S is a closed subset λ of S, which is a disjoint union of the simple geodesics. The geodesics of λ are called the *leaves* of λ. The lamination λ is called *minimal* if no proper subset of λ is a geodesic lamination. The following lemma gives an idea of the minimal laminations (our main object of study).

Proposition 1 *A minimal lamination λ in a closed orientable hyperbolic surface S is either a singleton (simple closed geodesic) or an uncountable nowhere dense (Cantor) subset of S.*

Proof. See Lemma 4.2.2 of [4] and Lemma 3.3 of [5]. □

The laminations λ, λ' on S are said to be (topologically) *equivalent* if there exists a homeomorphism $\varphi : S \to S$ such that each leaf of λ through the point $x \in S$ goes to the leaf of λ' through the point $\varphi(x)$. Clearly, the set of all laminations, $\Lambda(S)$, on S splits into the equivalence classes under the equivalence relation.

Since S is a complete hyperbolic surface, its universal cover is the unit disk D. Any $\lambda \in \Lambda(S)$ lifts to a lamination $\tilde{\lambda}$ on D, which is invariant under the action of the covering transformations. Every leaf of $\tilde{\lambda}$ is given by an unordered pair of points at the boundary of D, and therefore the space of geodesics is homeomorphic to a Möbius band, M.

Let $C(M)$ be the set of all closed subsets of M. The *Chabauty topology* on $C(M)$ is given by the (Hausdorff) distance d:

$$d(X,Y) \le \varepsilon \text{ iff } X \subseteq N_\varepsilon(Y) \text{ and } Y \subseteq N_\varepsilon(X), \tag{2}$$

where $N_\varepsilon(X)$ ($N_\varepsilon(Y)$) is a ε-neighbourhood of the closed set $X \in C(M)$ ($Y \in C(M)$). The function d turns the set $\tilde{\Lambda}$ of the laminations in D into a compact metrizable Hausdorff space. The Chabauty topology on $\Lambda(S)$ can be defined as a factor-topology of the topology on $\tilde{\Lambda}$ under the covering map. Everywhere in below, the standard topology on the set $\Lambda(S)$ will be the Chabauty topology. The following statement, mentioned in the introduction, will be critical.

Proposition 2 *The subspace $\Lambda^*(S)$ made of the singletons (i.e. the laminations consisting of one simple closed geodesic) is dense in the $\Lambda(S)$.*

Proof. See [4], Lemma 4.2.15. □

2.2 The principal regions and boundary leaves

If $\lambda \in \Lambda(S)$, then a component of $S - \lambda$ is called a *principal (complementary) region* for λ. (Note that $S - \lambda$ may have several connected components.) The leaves of λ, which form the boundary of a principal region, are called the *boundary leaves.* If λ is minimal (which we always assume to be the case), then each boundary leaf is a dense leaf of λ, isolated from one side. Note that by the proposition 1, it holds:

$$Area\ (S - \lambda) = Area\ S, \tag{3}$$

and therefore the principal region is a complete hyperbolic surface of the area $-2\pi\chi(S)$, where $\chi(S) = 2 - 2g$ is the Euler characteristic of the surface S. If U is a component of the preimage of the principal region in D, then U is a union of the ideal polygons U_i in D (see Fig.1)



(i) Maximal number of ideal polygons

(ii) Minimal number of ideal polygons

Figure 1: The region $U = \sqcup\ U_i$ for the surface $g = 2$

The hyperbolic area of an ideal n-gon U_i is equal to $(n-2)\pi$, see [5]. Since

$$\sum Area\ U_i = (4g - 4)\pi, \tag{4}$$

the number of the ideal polygons in U is finite.

2.3 The singularity data

Roughly, the singularity data counts the number and shape of the ideal polygons U_i, which cover the principal region of λ. Since $Area\ U_i = (n-2)\pi$ and

$\sum$ *Area* $U_i = (4g-4)\pi$, there exists only a finite number of the choices for such a data (for a fixed genus g). Let $\Delta = (k_1, \dots, k_m)$ be a set of the positive integers and the half-integers, such that $\sum k_i = 2g-2$. To each ideal n-gon U_i, we assign a number k_i such that

$$k_i = \frac{n-2}{2}. \tag{5}$$

The reader can verify that the equation $\sum$ *Area* $U_i = (4g-4)\pi$ is equivalent to the equation $\sum k_i = 2g-2$. An unordered tuple $\Delta = (k_1, \dots, k_m)$ will be called a *singularity data* of the lamination $\lambda \in \Lambda(S)$.

3 Proof of theorem 1

In this section the Legendre laminations are introduced. A slope of such laminations is defined and its properties are studied. Our theorem will follow from the properties and the singularity data. The proof is arranged into a series of lemmas.

An idea of the slope has been sketched in the introduction. Let us give an exact definition of a Legendre lamination and the associated slope. Recall that an n-th iterate of the regular continued fraction can be written as

$$\tau_0\tau_1\dots\tau_n(z) = p_0 + \cfrac{1}{p_1 + \cfrac{1}{p_2 + \dots + \cfrac{1}{p_n + z}}}, \tag{6}$$

where $p_0 \in \mathbb{N} \cup \{0\}$ and $p_k \in \mathbb{N}$ for $k = 1, \dots, n$, see the appendix. The Möbius transformations τ_k have the form:

$$\tau_0(z) = p_0 + z, \quad \tau_k = \frac{1}{p_k + z}, \quad k = 1, \dots, n. \tag{7}$$

One can rewrite the above transformations in the matrix form:

$$\tau_0 = \begin{pmatrix} 1 & p_0 \\ 0 & 1 \end{pmatrix}, \quad \tau_k = \begin{pmatrix} 0 & 1 \\ 1 & p_k \end{pmatrix}, \quad k = 1, \dots, n. \tag{8}$$

Let us call the Möbius transformation $\tau_k = \begin{pmatrix} 0 & 1 \\ 1 & p_k \end{pmatrix}$ a *regular factor* and the transformation $\tau_0 = \begin{pmatrix} 1 & p_0 \\ 0 & 1 \end{pmatrix}$ a *shift factor*. Let $\lambda^* \in \Lambda^*(X(N))$ ($\in \Lambda^*(X_0(N))$) be a simple closed geodesic. We denote by $(\lambda^*) \in \Gamma(N)$ ($\in \Gamma_0(N)$) an isometry, whose axis is λ^*, see the appendix.

Definition 1 *The geodesic lamination $\lambda \in \Lambda(X(N))$ ($\lambda \in \Lambda(X_0(N))$) is called Legendre if there exists a sequence of the singletons $\{\lambda_n^*\}$ convergent to λ, such that:*

$$(\lambda_0^*) \quad = \quad \tau_0$$

$$\begin{array}{rcl} (\lambda_1^*) & = & \tau_0\tau_1 \\ & \vdots & \\ (\lambda_n^*) & = & \tau_0\tau_1\dots\tau_n \\ & \vdots & \end{array} \tag{9}$$

where τ_0 is a shift factor and τ_n $(n = 1, \dots, \infty)$ are the regular factors.

Lemma 1 *The set of the Legendre laminations is a non-empty uncountable subset of $\Lambda(X(N))$ $(\Lambda(X_0(N)))$.*

Proof. First, let us evaluate the lengths of the closed geodesics approximating the Legendre lamination. A matrix calculus gives us:

$$\begin{array}{rcl} tr\ (\lambda_0^*) & = & 2 \\ tr\ (\lambda_1^*) & = & p_0 + p_1 \\ tr\ (\lambda_2^*) & = & 2 + p_0p_1 + p_1p_2 \\ tr\ (\lambda_3^*) & = & p_0 + p_1 + p_2 + p_3 + p_0p_1p_2 + p_1p_2p_3 \\ & \vdots & \end{array} \tag{10}$$

where $p_0 \in \mathbb{N} \cup \{0\}$ and $p_i \in \mathbb{N}$. Thus, the traces of (λ_n^*) is an unbounded monotone growing function of n. In view of the length-trace formula (see the appendix), we conclude that the lengths of the closed geodesics λ_n^* is also an unbounded monotone growing function of n. Since $\Lambda(X(N))$ $(\Lambda(X_0(N)))$ is compact, the sequence of the singletons $\{\lambda_n^*\}$ has a limit $\lambda \in \Lambda(X(N))$ $(\in \Lambda(X_0(N)))$. In view of the construction, the λ is a Legendre lamination. Thus, the Legendre laminations is a non-empty subset of the set $\Lambda(X(N))$ $(\Lambda(X_0(N)))$.

Let us prove that the set of the Legendre laminations is an uncountable set. Indeed, the set of the infinite sequences of the natural numbers p_k is an uncountable set. Since the subgroup $\Gamma(N)$ $(\Gamma_0(N))$ has a finite index in $SL(2, \mathbb{Z})$, there will be an uncountable set of the infinite sequences of p_k such that $(\lambda_n^*) \in \Gamma(N)$ $(\in \Gamma_0(N))$. The above sequences are bijective with the Legendre laminations. $\square$

In what follows, we shall work with the Legendre laminations. If it is not stated otherwise, by a lamination we shall mean the Legendre lamination.

Lemma 2 *If $\lambda \in \Lambda(X(N))$ $(\in \Lambda(X_0(N)))$ is a Legendre lamination, then the representation of λ by the sequence $\{p_i\}_{n=0}^{\infty}$ is unique.*

Proof. First, let us establish a bijection between $\{p_i\}_{i=0}^{n}$ and the positive rationals $\mathbb{Q}^+$ (and henceforth between the singletons λ_n^* and $\mathbb{Q}^+$). For that, let

$$r_n = p_0 + \cfrac{1}{p_1 + \cfrac{1}{p_2 + \dots + \cfrac{1}{p_n}}} \tag{11}$$

be the rational corresponding to $\{p_i\}_{i=0}^n$. Conversely, let us assume that n is even. Then the representation of any $r \in \mathbb{Q}^+$ by a regular continued fraction is unique, see [12]. Therefore we established a mapping $\Lambda^* \leftrightarrow \mathbb{Q}^+$, where Λ^* is the set of singletons.

Let to the contrary, $\{\lambda_n^*\}$ and $\{\bar{\lambda}_n^*\}$ be the two sequences of the singletons convergent to λ. Then, corresponding to $\{\lambda_n^*\}$ and $\{\bar{\lambda}_n^*\}$, there are two regular continued fractions convergent to the same limit. [1] This cannot happen in virtue of the proposition 4 (see the appendix). $\square$

Now one can define a slope of the Legendre lamination $\lambda \in \Lambda(X(N))$ ($\in \Lambda(X_0(N))$).

Definition 2 *Let $\lambda \in \Lambda(X(N))$ ($\lambda \in \Lambda(X_0(N))$) be a Legendre geodesic lamination whose regular factor decomposition is $\lambda = \lim_{n\to\infty} \tau_0\tau_1\dots$, where $\tau_0 = \begin{pmatrix} 1 & p_0 \\ 0 & 1 \end{pmatrix}$ and $\tau_n = \begin{pmatrix} 0 & 1 \\ 1 & p_n \end{pmatrix}$. Then the regular continued fraction*

$$\theta(\lambda) = p_0 + \cfrac{1}{p_1 + \cfrac{1}{p_2 + \dots}} \tag{12}$$

is called a slope of the lamination λ on the modular surface $X(N)$ ($X_0(N)$).

It may seem that the slopes are defined only for the laminations on the modular surfaces $X(N)$ and $X_0(N)$. It isn't true, because of the following construction, see [15], Chapter 8.5. Let $h : S_1 \to S_2$ be a homeomorphism between the Riemann surfaces S_1, S_2. Then it lifts to a homeomorphism $\tilde{h} : \mathbb{H} \to \mathbb{H}$ and extends to the boundary M of $\mathbb{H}$. Recall that the geodesic lamination is a closed subset of M. Thus, the mapping $\tilde{h} : C(M) \to C(M)$ sends the lamination $\lambda_1 \in \Lambda(S_1)$ to a lamination $\lambda_2 \in \Lambda(S_2)$, see [15], p.189 for the details. In particular, one has the following basic property of the slopes.

Lemma 3 *The slopes θ, θ' of the topologically equivalent laminations $\lambda, \lambda' \in \Lambda(X(N))$ ($\in \Lambda(X_0(N))$) are the equivalent irrationals, i.e. $\theta' = \theta \bmod GL(2, \mathbb{Z})$.*

Proof. Let λ have the regular factor decomposition

$$\lambda = \lim_{n\to\infty} \tau_0\tau_1\dots\tau_n\dots, \tag{13}$$

and let $\varphi : X(N) \to X(N)$ ($\varphi : X_0(N) \to X_0(N)$) be a homeomorphism sending λ to λ'. We want to find a regular factor decomposition of the λ'. If Λ^* is the set of singletons, then let $\varphi^* : \Lambda^* \to \Lambda^*$ be the action of φ on Λ^*. We established earlier a bijection $\Lambda^* \leftrightarrow \mathbb{Q}^+$. Thus, the action of φ extends to the positive rationals. But any such action has the form:

$$\varphi(r) = \frac{Ar+B}{Cr+D}, \quad r \in \mathbb{Q}^+, \quad \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in GL(2, \mathbb{Z}^+). \tag{14}$$

---

[1] Indeed, given natural order on the set $\{\lambda_n^*\} \cup \{\bar{\lambda}_n^*\}$, one can arrange this set into a Cauchy sequence (in the Chabauty topology). Such a sequence has a unique limit.

We shall use the Euclidean algorithm to decompose $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ into the regular factors $\tau'_0, \ldots, \tau'_k$, assuming that k is even. Therefore, the required regular factor decomposition of λ' has the form:

$$\lambda' = \lim_{n\to\infty} \tau'_0 \ldots \tau'_k \tau_0 \tau_1 \ldots \tau_n \ldots \tag{15}$$

Since the regular factor decompositions of λ and λ' have the same tails, we conclude in virtue of the proposition 5, that the slopes θ, θ' are the equivalent irrationals. $\square$

In view of the above, the mapping $\Lambda(X(N)) \to (\Theta, \Delta)$ $(\Lambda(X_0(N)) \to (\Theta, \Delta))$ is defined. Let us establish an inverse mapping. Let (Θ, Δ) be a pair consisting of the equivalence class Θ of the positive irrationals and $\Delta = (k_1, \ldots, k_m)$ the singularity data. First, we wish to construct a principal region $X(N) - \lambda$ $(X_0(N) - \lambda)$ of some lamination λ. It will be clear from the construction, that such a region defines the lamination λ itself.

By the definition, the principal region is given by the vertices of the ideal n-gons lying at the boundary, T, of the unit disk D. The total number d of such vertices depends on the singularity data $\Delta = (k_1, \ldots, k_m)$ and can be calculated according to the formula:

$$d = \sum_{i=1}^{m} (2k_i + 2). \tag{16}$$

(For example, the singularity data $\Delta = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ corresponding to the case (i) of Fig.1 will give $d = 12$, which is the maximal possible value for the genus $g = 2$. Similarly, the singularity data $\Delta = (2)$ corresponding to the case (ii) of Fig.1 will give $d = 6$, etc.) Thus, the principal region $U = U(\Delta)$ is specified by a point $p \in T^d$.

Let now $\theta \in \Theta$ be a positive irrational in the equivalence class Θ. We have shown earlier how to get an infinite sequence of the singletons $\lambda^*_n \in \Lambda^*(X(N))$ $(\lambda^*_n \in \Lambda^*(X_0(N)))$ out of the regular continued fraction of θ.

Let $\Omega_0 = \Omega_0(\lambda^*_0)$ be a subset of T^d consisting of the principal regions whose closure is disjoint from λ^*_0. Similarly, denote by $\Omega_1 = \Omega_1(\lambda^*_1)$ a subset of Ω_0 consisting of the principal regions, whose closure is disjoint from λ^*_1. In general, $\Omega_n = \Omega_n(\lambda^*_n)$ is a subset of Ω_{n-1} containing the principal regions, which do not intersect with the singleton λ^*_n. Thus one gets the following chain of the strict inclusions:

$$T^d \supset \Omega_0 \supset \Omega_1 \supset \ldots \supset \Omega_n \supset \ldots \tag{17}$$

It is not hard to see that the intersection $\cap_{i=0}^{\infty} \Omega_i$ consists of a single point $p \in T^d$. The point p defines a principal region $U = X(N) - \lambda$ $(U = X_0(N) - \lambda)$ and therefore a lamination $\lambda \in \Lambda(X(N))$ $(\lambda \in \Lambda(X_0(N)))$. By the construction, λ has the slope θ and the singularity data $\Delta = (k_1, \ldots, k_m)$. $\square$

4 Appendix

This section contains some basic facts on the continued fractions and the modular surfaces, which have been used in the proof of theorem 1. The reader is referred to [9] and [12] for a systematic account.

4.1 Continued fractions

The continued fraction can be viewed as a device to study the iterations of the Möbius transformations:

$$\tau_n(z) = \frac{a_n z + b_n}{c_n z + d_n}, \qquad z \in \mathbb{C}, \tag{18}$$

where $\{a_n\}$, $\{b_n\}$, $\{c_n\}$ and $\{d_n\}$ are given sequences of the complex numbers. The n-th iterate can be written as:

$$\tau_0 \tau_1 \dots \tau_n(z) = p_0 + \cfrac{q_0}{p_1 + \cfrac{q_1}{p_2 + \dots + \cfrac{q_{n-1}}{p_n + z}}}, \tag{19}$$

where $p_i, q_i \in \mathbb{C}$ are the rational functions of a_i, b_i, c_i and d_i, see [16]. With no restriction, one can study the iterations of the point $z = \infty$ since any $z_0 \in \mathbb{C}$ maps to the infinity under an iteration $\tau(z) = \frac{1}{z - z_0}$. In the case when the limit

$$\lim_{n \to \infty} \tau_0 \tau_1 \dots \tau_n(\infty) = v \tag{20}$$

exists and is finite, the continued fraction:

$$p_0 + \cfrac{q_0}{p_1 + \cfrac{q_1}{p_2 + \dots}} \tag{21}$$

is said to *converge* to the complex number v. Denote by T_n the n-th iteration of z:

$$T_n(z) = \tau_0 \tau_1 \dots \tau_n(z) = \frac{A_n z + B_n}{C_n z + D_n}. \tag{22}$$

The numbers p_n and q_n are called *n-th partial numerator* and *denominator* of the continued fraction, and the number $T_n(\infty) = \frac{A_n}{C_n}$ is called an *n-th approximant.* The numbers A_i, B_i, C_i and D_i depend on the partial numerators and denominators via the formulas:

$$\begin{array}{lll}
A_0 & = & p_0, \quad B_0 = q_0, \quad C_0 = 0, \quad D_0 = 1, \\
A_n & = & p_n C_{n-1} + q_n A_{n-1}, \qquad n = 1, 2, 3, \dots \\
B_n & = & p_n D_{n-1} + q_n B_{n-1}, \qquad n = 1, 2, 3, \dots \\
C_n & = & A_{n-1}, \quad D_n = B_{n-1}, \quad n = 1, 2, 3, \dots
\end{array} \tag{23}$$

Conversely, given the sequence of $\{A_i\}$, $\{B_i\}$, $\{C_i\}$ and $\{D_i\}$, one can recover the sequence of $\{a_i\}$, $\{b_i\}$, $\{c_i\}$ and $\{d_i\}$ (and, therefore, $\{p_i\}$ and $\{q_i\}$). The following statement is true.

Proposition 3 *Corresponding to a given system of the linear transformations*

$$T_n(z) = \frac{A_n z + B_n}{C_n z + D_n}, \quad A_n D_n - B_n C_n \neq 1 \tag{24}$$

there exists a uniquely determined sequence

$$\tau_n(z) = \frac{a_n z + b_n}{c_n z + d_n}, \quad a_n d_n - b_n c_n \neq 1 \tag{25}$$

such that $T_n(z) = \tau_0 \tau_1 \dots \tau_n(z)$.

Proof. See [8] p.28-29 for the proof, where also the explicit formulas for τ_n are given. □

In view of the proposition 3, a representation of the complex number v by a continued fraction is not unique. Indeed, any given sequence of the complex numbers $v_n \to v$ generates a continued fraction convergent to v. Note, that there are many distinct sequences $\{v_n\}$ convergent to v. However, the representation is unique over the field $\mathbb{Q}$. (This is the main reason behind our favor to the groups $\Gamma(N)$ $(\Gamma_0(N))$, as opposed to the general Fuchsian groups.) The continued fraction of the form

$$p_0 + \cfrac{1}{p_1 + \cfrac{1}{p_2 + \dots}} \tag{26}$$

is called a *regular continued fraction*, if all p_i are the positive integers except, may be, p_0 which can be 0. The regular continued fraction converges to a real number, which is a rational number if the fraction is finite, or an irrational number if the fraction is infinite. Recall that the regular continued fractions $\{p_n\}, \{p'_n\}$ are said to be distinct, if they differ in at least one term. The following is a well-known result.

Proposition 4 (Legendre) *Every positive irrational number has a unique decomposition into an infinite regular continued fraction.*

Proof. See e.g. [12] Satz 2.6, p. 33. □

The two irrationals θ, θ' are called *equivalent* if $\theta' = \theta \ mod \ GL(2, \mathbb{Z})$. By a *tail* of the infinite continued fraction $\{p_n\}_{n=0}^{\infty}$ one understands the sequence $\{p_n\}_{n=N}^{\infty}$, where $N \geq 0$. The continued fractions $\{p_n\}, \{p'_n\}$ are said to have the same tails, if $\{p_n\}_{n=N}^{\infty}$ is identical with $\{p'_n\}_{n=N'}^{\infty}$ for some positive integers N, N'. The following is a main property of the regular continued fractions.

Proposition 5 *The irrationals* θ, θ' *are equivalent if and only if their regular continued fractions have the same tails.*

Proof. See e.g. [12] Satz 2.24, p. 55. □

4.2 Modular surfaces

In this section the Riemann surfaces $X(N)$ and $X_0(N)$ are reviewed. The Riemann surfaces are routinely identified with the surfaces endowed with a hyperbolic metric (the uniformization theorem). We recommend Ch. 9, 11 of [9] for an extended exposition.

4.2.1 The principal congruence and Hecke subgroups

For an integer $N \geq 1$ let:

$$\begin{aligned} \Gamma(N) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2,\mathbb{Z}) \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \ mod\ N \right\} \\ \Gamma_0(N) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2,\mathbb{Z}) \mid c \equiv 0 \ mod\ N \right\}. \end{aligned} \tag{27}$$

The normal subgroups $\Gamma(N)$ and $\Gamma_0(N)$ have a finite index in the modular group $SL(2,\mathbb{Z})$ and are called the *principal congruence* and *Hecke subgroups* of the level N, respectively. The $\Gamma(N)$ and $\Gamma_0(N)$ consist of the matrices of the form:

$$\begin{pmatrix} 1+pN & qN \\ rN & 1+sN \end{pmatrix} \text{ and } \begin{pmatrix} p & q \\ rN & s \end{pmatrix}, \quad p,q,r,s \in \mathbb{Z}, \tag{28}$$

respectively. Clearly, $\Gamma(N) \subseteq \Gamma_0(N) \subseteq SL(2,\mathbb{Z})$.

4.2.2 The genus formulae

Let $\mathbb{H} = \{z = x + iy \in \mathbb{C} | y > 0\}$ be a Lobachevsky plane endowed with the hyperbolic metric $ds = |dz|/y$. The group $\Gamma(N)$ ($\Gamma_0(N)$) acts on $\mathbb{H}$ by the linear transformations $z \mapsto \frac{az+b}{cz+d}$, where $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma(N) (\in \Gamma_0(N))$. The left coset $\Gamma(N)\backslash\mathbb{H}$ ($\Gamma_0(N)\backslash\mathbb{H}$) is an orbifold with the cusps corresponding to the stabilizers of the group $\Gamma(N)$ ($\Gamma_0(N)$). We wish to compactify the above orbifold to a Riemann surface. Let $\mathbb{Q}^* = \mathbb{Q} \cup \{\infty\}$. One can add a boundary line to $\mathbb{H}$, so that $\mathbb{H}^* = \mathbb{H} \cup \mathbb{Q}^*$. $\mathbb{H}^*$ is the Hausdorff space with a discrete action of $SL(2,\mathbb{Z})$, see [9] p. 311. Then

$$X(N) = \Gamma(N)\backslash\mathbb{H}^* \quad \text{and} \quad X_0(N) = \Gamma_0(N)\backslash\mathbb{H}^* \tag{29}$$

are the compact Riemann surfaces. The $X(N)$ and $X_0(N)$ are called the *modular (Riemann) surfaces* of the level N. The genus of $X(N)$ is given by the formula:

$$g = 1 + \frac{1}{24}\tau N^2(N-6)\prod_{p|N}\left(1 - \frac{1}{p^2}\right), \qquad \tau = \begin{cases} 2, & \text{if } N = 2, \\ 1, & \text{if } N > 2 \end{cases}, \tag{30}$$

see [7], p.15. An explicit formula for the genus of $X_0(N)$ is slightly longer:

$$g = 1 + \frac{\mu(N)}{12} - \frac{\mu_2(N)}{4} - \frac{\mu_3(N)}{3} - \frac{\mu_\infty(N)}{2},$$

$$
\begin{aligned}
\mu(N) &= N\prod_{p|N}(1+\frac{1}{p}),\\
\mu_2(N) &= \begin{cases} 0, \text{ if } 4|N, \\ \prod_{p|N}\left(1+\left(\frac{-1}{p}\right)\right), \text{ otherwise}\end{cases}\\
\mu_3(N) &= \begin{cases} 0, \text{ if } 2|N \text{ or } 9|N \\ \prod_{p|N}\left(1+\left(\frac{-3}{p}\right)\right), \text{ otherwise}\end{cases}\\
\mu_\infty(N) &= \sum_{d|N}\varphi(GCD\ (d, N|d)),
\end{aligned}
\tag{31}
$$

where $\left(\frac{-1}{p}\right)$, $\left(\frac{-3}{p}\right)$ are the Legendre symbols and φ is the Euler function, see [9], Theorem 9.10. It is well known, that there exist (a finite number of) the modular surfaces of any given genus $g \geq 0$.

4.2.3 The length-trace formula

An element $\tau \in SL(2, \mathbb{Z})$ is called:

(i) elliptic if $|tr\ (\tau)| < 2$;

(ii) parabolic if $|tr\ (\tau)| = 2$;

(iii) hyperbolic if $|tr\ (\tau)| > 2$.

The cases (i)-(iii) correspond to the isometries of $\mathbb{H}$ with: (i) one fixed point in the interior of $\mathbb{H}$, (ii) one fixed point at the boundary of $\mathbb{H}$, (iii) two fixed points at the boundary of $\mathbb{H}$. If $\tau \in \Gamma(N)$ $(\tau \in \Gamma_0(N))$ is a hyperbolic isometry with the two fixed points $a \neq b$ at the boundary of $\mathbb{H}$, let $\tilde{\lambda} = \tilde{\lambda}(a, b)$ be an open geodesic arc through a and b. If $z \in \tilde{\lambda}$, then $\tau(z) \in \tilde{\lambda}$ and the length

$$
l([x, \tau(x)]) = 2\ cosh^{-1}\left(\frac{1}{2}|tr\ \tau|\right), \tag{32}
$$

of the hyperbolic arc $[z, \tau(z)]$ is independent of z, see [7]. On the other hand, τ is a covering isometry. Therefore, the arc $[z, \tau(z)]$ covers a closed geodesic $\lambda \in \Lambda(X(N))$ $(\lambda \in \Lambda(X_0(N)))$ of the length $l([z, \tau(z)])$.

Proposition 6 *Each simple closed geodesic* $\lambda \in \Lambda(X(N))$ $(\lambda \in \Lambda(X_0(N)))$ *comes from an equivalence class* $[\tau] = \{g\tau g^{-1} \mid g \in SL(2, \mathbb{Z}\}$ *of a hyperbolic transformation* $\tau \in \Gamma(N)$ $(\tau \in \Gamma_0(N))$.

Proof. See e.g. [7]. □

The geodesic λ in the proposition 6 is called an *axis* of the hyperbolic transformation τ.

References

[1] D. V. Anosov, Flows on closed surfaces and behavior of trajectories lifted to the universal covering plane, J. of Dynamical and Control Syst. 1 (1995), 125-138.

[2] D. V. Anosov and E. V. Zhuzhoma, Nonlocal asymptotic behavior of curves and leaves of laminations on universal coverings, Proceedings of the Steklov Institute of Mathematics ; v. 249, 2005, issue 2.

[3] E. Artin, Ein mechanisches System mit quasiergodischen Bahnen, Abh. Math. Sem. Hamburg (1924), 170-175; in: Collected papers, 499-504, Springer Verlag.

[4] R. D. Canary, D. B. A. Epstein and P. Green, Notes on notes of Thurston, Lond. Math. Soc. Lect. Notes 111 (1987), 3-92.

[5] A. J. Casson and S. A. Bleiler, Automorphisms of Surfaces after Nielsen and Thurston, Lond. Math. Soc. Student Texts 9, Cambridge, 1988.

[6] A. Fathi, F. Laudenbach and Z. Poénaru, Travaux de Thurston sur les surfaces, Séminaire Orsay, France, Astérisque 66-67 (1979).

[7] R. C. Gunning, Lectures on Modular Forms, Princeton Univ. Press, 1962.

[8] W. B. Jones and W. J. Thron, Continued Fractions: Analytic Theory and Applications, Encyclopedia of mathematics and its applications, Vol. 11, Addison-Wesley 1980.

[9] A. W. Knapp, Elliptic Curves, Princeton Math. Notes, 1992.

[10] R. Moeckel, Geodesics on modular surfaces and continued fractions, Ergod. Th. & Dynam. Sys. 2 (1982), 69-83.

[11] P. J. Myrberg, Ein Approximationssatz für die Fuchsschen Gruppen, Acta Math. 57 (1931), 389-409.

[12] O. Perron, Die Lehre von den Kettenbrüchen, Bd.1, Teubner, 1954.

[13] S. Schwartzman, Asymptotic cycles, Annals of Math. 66 (1957), 270-284.

[14] C. Series, The modular surface and continued fractions, J. London Math. Soc. 31 (1985), 69-80.

[15] W. P. Thurston, The Geometry and Topology of Three-Manifolds, MSRI 1997, electronic edition of 1980 Princeton Univ. notes, available at http://www.msri.org/gt3m/

[16] H. S. Wall, Analytic theory of continued fractions, Chelsea, N.Y., 1973

[17] A. Weil, Les familles de courbes sur le tore, Mat. Sbornik 1(1936), 779-781.

The Fields Institute for Mathematical Sciences, Toronto, ON, Canada, E-mail: igor.v.nikolaev@gmail.com